

The value of oceanic marine reserves for protecting highly mobile pelagic species: Coral Sea case study



Daniela M. Ceccarelli

Jan. 31, 2011

Report prepared for the Protect Our Coral Sea campaign



Contents

| | |
|---|----|
| Executive summary..... | 3 |
| Introduction | 4 |
| Positive effects of marine reserves for pelagic and migratory species | 5 |
| Positive ecosystem-level effects of pelagic marine reserves | 10 |
| Predictability of bathymetric and hydrographic features | 11 |
| Key arguments from marine reserve critics..... | 11 |
| Argument 1: Reserve size and migration distance..... | 11 |
| Argument 2: Fisheries losses and overexploitation | 15 |
| Argument 3: Governance and enforcement..... | 18 |
| Parallels with Chagos/BIOT..... | 22 |
| Conclusions | 22 |
| Acknowledgments..... | 23 |
| References | 24 |
| Appendix 1—Migratory and pelagic species of the Coral Sea..... | 35 |

Figures and tables

| | |
|---|----|
| Table 1. Summary of studies predicting marine reserve benefits..... | 6 |
| Table 2. Recorded movement distances for species caught by the ETBF | 13 |
| Figure 1. Landings by functional groups on the northeast Australian shelf..... | 17 |
| Table 3. Catches (tonnes) of the most important ETBF species between 1998 and 2006..... | 17 |
| Table 4. Summary table of challenges raised by critics of marine reserves..... | 19 |

Executive summary

This paper reviews the value of marine reserves for protecting migratory and highly mobile pelagic species. Three common arguments used by critics of marine reserves were considered, and the proposed Coral Sea marine reserve was highlighted as an example of how these challenges could be met. Comparisons were made with the UK's recently established large-scale no-take marine reserve in the Chagos Archipelago. The major findings of this review are as follows:

- Recent scientific findings support the idea that large pelagic species benefit from marine reserves, because:
 - o Protecting even a part of species' range or life cycles, especially critical habitat areas which function as important feeding or breeding grounds, reduces overall population mortality. Partial protection works best in a coordinated approach for protection, including no-take marine reserves and areas of limited fishing effort;
 - o Pelagic species are not uniformly distributed, but tend to aggregate around bathymetric and hydrographic features that are predictable in space and time, making the most beneficial design of pelagic reserves possible; and
 - o Even species identified as highly migratory display predictable movement patterns where the majority (70-90 percent) of the population moves no farther than 600 kilometres (km).
- Critics of marine reserves for the conservation of pelagic species have focused on three major issues:
 - o Size: Reserves have to be very large to encompass an adequate portion of pelagic species' home ranges;
 - o Fisheries management: There is some scepticism about the use of marine reserves as a fisheries management tool, with concerns about displaced fishing effort; and
 - o Governance: There is concern that arranging governance and enforcement beyond one country's exclusive economic zone (EEZ) is too challenging.
- The proposed Coral Sea marine reserve is ideal to surmount these challenges and take advantage of the benefits, in the following ways:
 - o Size: The proposed reserve encompasses almost 1 million km², making it large enough to protect a significant portion (if not all) the home ranges and life cycles of many pelagic species that reside within it. It is large enough to also encompass a large variety of bathymetric and hydrographic features that provide key habitat for pelagic species at vulnerable times (feeding and breeding).
 - o Fisheries management: The Coral Sea marine reserve is being proposed for broad ecosystem conservation benefits, not as a fisheries management tool. Its history of relatively low exploitation by fisheries means that pelagic, reef and deep-sea ecosystems are relatively undisturbed, making them strong candidates for conservation to prevent future exploitation, and displaced fishing would be minimal. The permanent closure of the Coral Sea would partially or wholly protect populations of species that are subject to overexploitation elsewhere along their range.

- Governance: The proposed Coral Sea marine reserve lies entirely within Australia's EEZ, making negotiations with neighbouring countries unnecessary. Precedents for managing large marine reserves exist in Australia, and the proposed reserve design (a single large no-take area) is the most cost-effective management option.

Introduction

The open ocean, long considered invulnerable, is under increasing pressure from human impacts^{1,2}. As coastal fisheries become depleted, and technological improvements allow fishing vessels to venture farther offshore, large migratory marine species have become more intensely exploited^{3,4}. More than 10 years ago, researchers reported that “almost 70 percent of fished stocks are listed as ‘fully fished, overfished, depleted, or recovering’ ”⁵. Numerous heavily exploited species are now of conservation concern, including tuna, billfish and sharks^{3,6-8}. For instance, almost all sharks recorded by Baum *et al.* (2003)³ underwent a 50 percent decline over the 15 years of their study. While marine reserves have steadily increased in coastal areas⁹, there is still a lack of adequate protection for pelagic ecosystems worldwide^{1,4}.

Marine reserves are considered the best conservation tools available to protect marine species and habitats from exploitation¹⁰. However, the global percentage of area protected in marine reserves is less than 1 percent¹¹, well below the proportion needed to adequately represent all biogeographic zones^{5,12}. Ideally, marine reserves would maintain or restore native species diversity, habitat diversity and heterogeneity, keystone species and connectivity¹³. Diverse and complex habitats promote species diversity¹⁴⁻¹⁶, which in turn affects productivity¹⁷, resistance to and recovery from disturbance¹⁸, stable food web dynamics^{6,19} and the capacity to fill all the roles required for a functioning ecosystem even if individual species disappear²⁰. Keystone species, or species that can affect whole ecosystems through changes in their abundance^{6,21}, and connectivity between habitats and populations are also considered critical to the persistence of ecosystems in the face of stressors¹⁶. The benefit of no-take marine reserves to the abundance and size of sedentary species, especially those with a history of exploitation, is well-documented²²⁻²⁴. Despite gaps in knowledge and persisting doubts about their effectiveness in protecting pelagic and migratory species²⁵, increasing evidence suggests that marine reserves can also benefit these highly mobile species²⁶⁻²⁸.

A recent increase in the establishment of very large marine reserves that encompass pelagic and deep-sea ecosystems mirrors the positive shift in the perception of their usefulness²⁹. The proposed no-take Coral Sea marine reserve would be one of the world's largest, covering around 1 million km² between the seaward edge of the Great Barrier Reef Marine Park and the border of Australia's EEZ³⁰. The proposed reserve aims to protect geological, ecological, cultural and heritage values represented within the area³⁰. The Coral Sea is regionally important, providing breeding and spawning grounds for a number of migratory species, nesting habitat for seabirds and turtles, and dispersal stepping-stones for marine species between the Pacific and the Great Barrier Reef. There is some concern that this reserve would not adequately protect highly mobile and migratory pelagic species, but parallels exist with other large-scale marine reserves where the protection of pelagic species has been one of the key goals²⁷.

This paper reviews scientific research that assesses the benefits of marine reserves for protecting migratory and highly mobile pelagic species. This review also evaluates the case of the proposed Coral Sea marine reserve, and considers the three most common arguments against marine reserve establishment in the context of pelagic species protection. Because of similar issues, comparisons are made with the UK's recently established large no-take marine reserve in the Chagos Archipelago.

Positive effects of marine reserves for pelagic and migratory species

Positive, measurable marine reserve effects on pelagic species exist^{3,31-33}, including for large migratory species such as marine mammals and large predators^{32,34-36}. The protection of such species over their whole annual range may not be realistic, but marine reserves can be used to protect them at highly vulnerable stages of their life cycles by encompassing seasonal breeding or feeding grounds, or key parts of their migratory routes^{34,37-39}, all habitats critical to the survival of the species (critical habitat)⁴⁰. Because highly mobile species are often also the most heavily exploited, they may receive some benefit even from small marine reserves, although the probability of adequate protection increases with increasing reserve size²⁸. Spawning aggregations in particular are increasingly shown to be temporally predictable, and therefore both vulnerable to exploitation and responsive to protection^{23,36,41,42}. Protecting vulnerable areas such as breeding or spawning grounds can result in a greater source of larvae or young to the exploited part of the population, resulting in improved breeding success and lower mortality overall^{36,39,43}. Some researchers have suggested that protecting at least 50 percent of a species' total habitat would afford it adequate protection^{43,44}, while others argue that for species that undertake extensive migrations, spatial protection must be coupled with strict harvest quotas⁴⁵.

It is generally agreed that one large reserve is more effective for protecting wide-ranging species than a number of small reserves, even if they protect the same overall percentage of a region⁴⁶. A review of marine reserves in 1999 concluded that large migratory species could not be protected with small reserves, where the largest reserve measured 350 km²⁵. In contrast, a later review of marine reserve success included reserves in excess of 1,000 km² and found that populations of lobsters, snappers and other fish with large seasonal movements of up to 1,000 km⁴⁷ did benefit substantially, even if the reserves encompassed only a part of their home range²³.

Critics argue that marine reserves cannot benefit mobile commercial species and are only appropriate for small-scale fisheries in tropical regions^{23,48}. The argument against pelagic marine reserves is driven by the notion that they don't work as fisheries management tools; however, their capacity to protect ocean biodiversity is less disputed⁴⁹. Protection for pelagic species already exists in some places (see also Hooker and Gerber 2004³⁴), including:

- A number of temporary spatial closures or bans on specific gear types to protect pelagic and migratory stocks in the high seas by regional fisheries management organisations around the world⁵⁰;
- Three areas totalling 17,000 km² in the Georges Bank, Gulf of Maine, partially closed to fishing²³;
- The Indian Ocean and Southern Ocean Whale Sanctuaries⁵⁰,

- Temporary closures by the Mexican government of fixed areas to commercial longlining off the coast of Baja California³²;
- The seasonal protection of southern bluefin tuna habitat off eastern Australia⁵¹;
- Cetacean feeding, migration or calving areas in the Mediterranean Sea, the Great Australian Bight off southern Australia and the Southern Ocean^{34,45,52,53};
- Four areas that exclude certain fishing methods (especially purse seining) totalling 1.2 million km² stretching 7,000 km from French Polynesia to Palau, established by western Pacific island nations⁵⁴;
- The 356,893-km² Papahānaumokuākea Marine National Monument in the Northwestern Hawaiian Islands;
- The 246,000-km² Mariana Trench Marine National Monument in the northern Mariana Islands, with 17 percent closed to fishing; and
- The 544,000-km² Chagos Protected Area in the Indian Ocean²⁹.

The current lack of data from large oceanic marine reserves means that some of the evidence of their positive effects for pelagic species comes from modelling studies (Table 1)^{3,55}. However, the most compelling proof comes from what is termed an “unintended experiment”: Fisheries catch data were compiled after a large area of the North Sea (575,000 km²) had been closed to fishing during the six years of World War II³⁵. This study showed conclusively both increased abundance in pelagic species and larger proportions of older fish. In a more recent study, the protection of billfish from longlining in a part of their range off Baja California resulted in an overall population increase of up to 22 percent³². The recovery of whales following the combination of reduced whaling and the establishment of large whale sanctuaries also highlights the benefits of marine reserves that cover only parts of species’ ranges^{56,57}. More recent reviews and meta-analyses have found increasing empirical evidence that highly mobile and large-bodied species exploited by fisheries tend to benefit from marine reserves of varying size^{26,33,37}.

Table 1. Summary of studies predicting marine reserve benefits for pelagic and migratory species, using a variety of methods.

| Source (method) | Location | Species | Summary of findings |
|--|--------------|-----------------------------|---|
| Clark (1996) ⁴⁴ (discussion paper) | Global | All exploited species | - Marine reserves need to include up to 50% of a population or home range in order to protect a species from overfishing. |
| Lauck <i>et al.</i> (1998) ⁴³ (modelling) | Hypothetical | Hypothetical | - Marine reserves need to include up to 50% of a population or home range in order to protect a species from overfishing. |
| Roberts and Sargent (2002) ³⁶ | Hypothetical | Hypothetical migratory fish | - Protecting important aggregation areas has a disproportionate effect on |

| Source (method) | Location | Species | Summary of findings |
|--|----------------------|--------------------------------------|--|
| (modelling) | | | entire populations of highly mobile and migratory species. |
| Baum <i>et al.</i> (2003) ³ (modelling) | Northwest Atlantic | Sharks | <ul style="list-style-type: none"> - Priority areas for shark conservation are highlighted. - Population benefits for sharks with fishing closures of different areas are modelled. - Marine reserves coupled with reductions in fishing effort have positive effects on sharks and other large pelagic predators. |
| Gell and Roberts (2003) ²³ (review) | Global | All species | <ul style="list-style-type: none"> - Highlights reversal of notion that mobile species cannot be protected by marine reserves. - Even for highly mobile species, a portion of the population may remain within a small home range - Protecting migration bottlenecks, nurseries, spawning or feeding aggregation sites can benefit even highly migratory species. |
| Worm <i>et al.</i> (2003) ³¹ (modelling) | Northwest Atlantic | Pelagic species, primarily predators | <ul style="list-style-type: none"> - Identify pelagic diversity hot spots associated with productivity and habitat features. - Protecting hot spots from fishing has large benefits for pelagic populations. - Identify pelagic predator diversity hot spot in Great Barrier Reef/Coral Sea area. |
| Willis <i>et al.</i> (2003) ²⁶ (empirical, modelling) | Northern New Zealand | Snapper (<i>Pagrus auratus</i>) | <ul style="list-style-type: none"> - Density and size of snapper increase inside marine reserves, despite its high mobility. |
| Hooker and Gerber | Global | Predators and megafauna | <ul style="list-style-type: none"> - Marine reserves are beneficial for protecting predators and |

| Source (method) | Location | Species | Summary of findings |
|---|---------------------------|---|--|
| (2004) ³⁴ (discussion paper) | | | other megafauna (e.g., cetaceans, seabirds). - Present tools and approaches for enhancing marine reserve effectiveness. |
| Micheli <i>et al.</i> (2004) ²⁸ (meta-analysis) | Global | All species | - Highly mobile species benefit from marine reserve protection. |
| Palumbi (2004) ³⁷ (review) | Global | All species | - Is ambivalent about the value of marine reserves for migratory pelagic species but states that "If fishing effort is not displaced, then the impact of reserves on highly migratory species is similar to the effect of decreasing fishing effort by the same percentage as the percent area dedicated to reserves." |
| Hyrenbach <i>et al.</i> (2006) ⁵⁸ (empirical) | Central California | Black-footed albatross (<i>Phoebastria nigripes</i>) | - Advocates protecting albatross foraging grounds, even though these comprise only a part of their overall range. |
| Louzao <i>et al.</i> (2006) ³⁸ (empirical, modelling) | Balearic Islands | Balearic shearwater (<i>Puffinus mauretanicus</i>) | - Marine zoning measures can benefit populations of far-ranging seabirds by extending protective measures beyond their breeding colonies. |
| Alpine and Hobday (2007) ⁵⁵ (modelling) | Eastern Australia | Pelagic, migratory and other species targeted by fisheries or of conservation concern | - Quantified the area requirements of pelagic protected area networks to protect pelagic species (target and non-target). - Area requirements ranged from 7 to 26% of the region for adequate protection of pelagics. |
| Pichegru <i>et al.</i> (2009) ³⁹ (empirical) | Benguela upwelling region | Cape gannets (<i>Morus capensis</i>) and African | - Measured overlap between seabird feeding and commercial fishing grounds. |

| Source (method) | Location | Species | Summary of findings |
|---|--------------------------|--|---|
| | | penguins (<i>Spheniscus demersus</i>) | - Marine reserves in bird-feeding hot spots is likely to increase the birds' breeding success. |
| Beare <i>et al.</i> (2010) ³⁵ (empirical) | North Sea | North Sea gadoids | - Large North Sea area unfished during World War II. - Large benefits to exploited fish, including migratory species. - Older fish benefit fastest and in greatest proportion, creating a "Mexican wave" in numbers of subsequent generations. |
| Claudet <i>et al.</i> (2010) ³³ (meta-analysis) | European marine reserves | Fish | - Density and size of species targeted by fisheries increase inside marine reserves, even highly mobile species. |
| De Juan and Leonart (2010) ⁴ (modelling) | Mediterranean | All pelagic species | - Identifies habitats critical to pelagic species in the Mediterranean. - Advocates for marine reserve protection of pelagic species. |
| Jensen <i>et al.</i> (2010) ³² (empirical) | Baja California | Striped marlin (<i>Kajikia audax</i>) | - Temporary closures of Mexico's EEZ to long-lining (1977–1980, 1984–1985) caused increase in striped marlin, despite its range extending outside the closed area. |
| Koldewey <i>et al.</i> (2010) ²⁷ (review) | Global | All species | - Increasing evidence that even partial protection of highly mobile and migratory species is beneficial. - "Highly migratory" species may be based on long-range movements of a few individuals, while most of the population remains within a home range. |

The Coral Sea plays a regionally important role for South Pacific pelagic fish stocks. It hosts spawning aggregations of tuna, billfish⁵⁹⁻⁶¹ and other large pelagic fish⁶², and contains nesting grounds for marine turtles and seabirds^{63,64}. It is one of the few oceanic areas that has retained a high diversity of large oceanic predators³¹ that are heavily exploited elsewhere along their range⁶⁵. The wide range of bathymetric features such as reefs and seamounts attracts migratory species, including seabirds⁶⁶ and cetaceans⁶⁷. The proposed Coral Sea marine reserve, covering an area of around 1 million km², would ensure that a high proportion of the key areas in the life cycles and migration pathways of many pelagic species are protected (see Appendix 1).

Positive ecosystem-level effects of pelagic marine reserves

The capacity for well-designed marine reserves to meet the objectives of maintaining or restoring species, habitats and connectivity¹³ is increasingly well-documented, especially for shallow-water environments^{10,24}. When species diversity is protected, for instance, rates of resource collapse decrease and recovery potential, stability and water quality increase, sometimes exponentially¹⁷. Most scientists agree that marine reserves protect habitats and biodiversity^{68,69} by reducing stressors such as extraction⁵⁵, making ecosystems more resilient and therefore better able to withstand the more pervasive effects of climate change and pollution^{17,70-72}.

Because of the vast extent of oceanic ecosystems and the increased understanding of their value, it is now believed that pelagic marine reserves are rapidly becoming a reality in the planet's "last frontier of conservation management"^{11, 28,54,73}, although their implementation may be both costly and difficult⁷⁴. Large-scale pelagic marine reserves have the capacity to protect greater diversity, larger habitats and entire trophic webs and ecological processes^{29,34}. As well as pelagic habitats, they would include deep-sea ecosystems that are poorly understood and yet at risk from activities such as trawling, oil and mineral exploration, and sea dumping^{71,73,75}. Protecting entire bioregions has the benefit that pathways of connectivity will be preserved without the need for complex conservation planning to establish a marine reserve network¹⁶.

The proposed marine reserve in the Coral Sea would encompass four large-scale bioregions⁶², including terrestrial, pelagic, shallow, deep and abyssal ecosystems⁶². The cessation of fishing would not only benefit the numerous target species, but also the large number of species caught as bycatch⁷⁶. The coral reefs in the Coral Sea are already vulnerable to high temperature anomalies that can cause coral bleaching and death; some of these reefs appear more vulnerable than others⁷⁷. Studies of genetic connectivity^{78,79} indicate that the Coral Sea might contain the entire genetic stock of some species, as well as the stepping-stones of dispersal between the wider Pacific Ocean and the Great Barrier Reef⁸⁰. Preliminary research suggests that the Coral Sea's deep-sea habitats are vulnerable, patchy and important for deep-diving pelagic species⁸¹. A permanent closure would safeguard critical habitats such as breeding and feeding hot spots, support the resilience of ecosystems and connectivity pathways, and therefore enhance the persistence of pelagic populations.

Predictability of bathymetric and hydrographic features

Despite the perception that the open ocean is relatively featureless and large species move through it more or less at random, there are persistent and predictable bathymetric and hydrographic features^{3,31,82}. Hyrenbach *et al.* (2000)⁸² distinguished between three types of oceanic features that can be mapped: 1) static systems, defined by topographic features; 2) persistent hydrographic features, such as currents and frontal systems; and 3) ephemeral habitats, shaped by wind- or current-driven upwellings and eddies. All three types of features are known to attract aggregations of marine life and may be identified by analysing the foraging distribution of higher predators⁸² or by making use of sophisticated real-time imagery¹¹. The tendency for pelagic species to aggregate in predictable seasons and areas of ocean has made them highly exploitable, and predicting the location of aggregations of commercially valuable species has been important in fisheries oceanography⁵⁵. Even highly migratory species have been found to travel along predictable pathways defined by topographic or oceanographic features⁸³. In fact, the predictability of movements and aggregations of terrestrial migratory species has already been applied to the protection of migratory species on land and is considered viable in the open ocean⁷³³.

Topographic features such as shelf breaks, canyons and seamounts alter the water flow above them, causing highly productive plankton blooms or swarms⁸². Turbulent water flow in the lee of islands and emergent reefs also serves to retain the planktonic food sources of pelagic fish⁸⁴. These features have been shown to act as highly effective natural aggregation devices for tuna and other migratory species, primarily for feeding^{85,86}, but also for breeding⁸⁷. Hydrodynamic features such as eddies, currents, upwellings, downwellings or fronts are also areas of high productivity, attracting species from all trophic levels⁸². Temperature fronts, or water mass boundaries, are well-known as biological hot spots and migration corridors^{88,89}. The dynamic nature of these features has served to fuel the argument against spatial closures^{51,55}, but many of these features are predictable in space and time and can be tracked remotely¹¹.

The Coral Sea contains more than 30 emergent reefs and atolls, part of a major seamount chain, four major plateaus and a series of troughs, slopes, canyons and abyssal plains^{62,90}. Major currents, gyres and eddy systems interact with these bathymetric features, creating a complex pattern of hydrodynamic regimes with the potential to support numerous hot spots of pelagic diversity⁹¹. Most of these features and communities remain to be studied, but existing research has shown increased productivity in the lee of islands and reefs⁸⁴ and an increase in the abundance of the larvae of some tuna and billfish species near reefs and islands⁹². A further study has indicated that broadbill swordfish may set up resident populations in the vicinity of seamounts, and favourable conditions for this exist in the Coral Sea⁶⁰. Sophisticated remote sensing tools exist that can map the position, movement and extent of areas likely to attract aggregations of pelagic life¹¹. Protecting the entire Coral Sea would ensure the inclusion of these features, and the large size of the reserve would allow for seasonal or annual movements of the more dynamic fronts, eddies and currents.

Key arguments from marine reserve critics

Argument 1: Reserve size and migration distance

A key argument against oceanic marine reserves is that they would need to be exceedingly large to adequately protect migratory species⁷⁴. It is argued that even species that habitually

reproduce or feed in an area of ocean do not exhibit the level of site fidelity needed for a marine reserve. For instance, Kaplan *et al.* (2010)⁷⁴ argue that the tropical skipjack tuna (*Katsuwonus pelamis*) does not undertake consistent feeding or breeding migrations, making it difficult to establish a reserve for its protection in the right place. Similarly, the population of whale sharks protected by the Ningaloo Marine Park is in decline, due to intensifying threats elsewhere along its range⁹³. However, it has previously been shown that even protecting one part of a tuna's home range or life cycle can have an overarching positive effect on its overall stocks⁹⁴. A modelling study showed that because highly mobile species are often also the most heavily exploited, they are most likely to benefit from marine reserves, with larger reserves affording greater protection²⁸. Partial protection for migratory species cannot be considered futile^{4,95}.

In fisheries management, the phrase "highly migratory" can be derived from the long-range movements of a few individuals within a population, with studies of tuna mobility demonstrating they would benefit from national-level closures⁹⁴. For instance, in assessing the efficacy of the Chagos marine reserve for the protection of tuna stocks, Sheppard (2010)²² questioned the distances implied by "highly migratory" and found that the average distance travelled by tagged tunas meant that they would spend relatively little time outside the reserve. Even for a species shown to be highly migratory (the striped marlin [*Kajikia audax*]), reduction in fishing pressure over a small part of its range resulted in measurable improvement³². Another study cautions that long-range movement may be undertaken by only some members of a population and that generalizations about the mobility of the entire population may be inappropriate²⁶.

Applying this argument to the Coral Sea, migration distances reported in the available literature were collated for the pelagic species most targeted by the Eastern Tuna and Billfish Fishery, after Hobday (2010)⁹⁶. If movement data were available, further species were added if they were highly valued by the game-fishing industry or often caught as bycatch (Table 1). Species with the ability to undertake migrations of more than 1,000 nautical miles tend to move between 400-600 nautical miles on average (Table 2). While this may take them into the EEZs of other Pacific nations, there is a high likelihood that they may spend 50 percent of their time inside the Coral Sea, effectively complying with the guidelines set up by Clark (1996)⁴⁴ and Lauck *et al.* (1998)⁴³ for adequate protection. Sibert and Hampton (2003)⁹⁴ state that while international arrangements are ideal, protection of tuna stocks within an individual country's EEZ is also highly effective.

Table 2. Recorded movement distances for species caught by the Eastern Tuna and Billfish Fishery, ranked in order of importance after Hobday (2010)⁹⁶. Also added are five species (below the line) that feature strongly in fisheries, as target or bycatch species. For a more comprehensive list of pelagic and migratory species that are exploited in the Coral Sea or elsewhere, and may therefore benefit from protection, see Appendix 1.

| Species | Info | Reference |
|----------------------------|---|--|
| Yellowfin tuna | <p>34 tunas tagged in the Coral Sea and recaptured, most along the New South Wales (NSW) coast within 200 nautical miles (nm) of release; longest straight-line distance between release and recapture was 569 nm after 9 months.</p> <p>273 tunas tagged by game fishers, most recaptured within the Australian Fishing Zone less than 600 nm from release.</p> <p>Most tunas tagged by the Australian Commonwealth Scientific and Research Organization (CSIRO) in the Coral Sea were caught close to the release area.</p> | <p>Hampton and Gunn (1998)⁹⁷</p> |
| | <p>Median lifetime displacement of 336-376 nm, mostly northeast into EEZs of other Pacific Island nations.</p> | <p>Sibert and Hampton (2003)⁹⁴</p> |
| | <p>Average distance travelled for all individuals recaptured from 2006 to 2008 was 247 nm.</p> | <p>Industry & Investment (I&I) NSW (2009)⁹⁸</p> |
| Broadbill swordfish | <p>Median movement of tagged fish was 744 km.</p> | <p>Sedberry and Loefer (2001)⁹⁹</p> |
| | <p>Average distance travelled during 193 days was 30 ± 43 km.</p> | <p>Sepulveda <i>et al.</i> (2010)¹⁰⁰</p> |
| Bigeye tuna | <p>Most tuna tagged by CSIRO in the Coral Sea were caught close to the release area.</p> | <p>Hampton and Gunn (1998)⁹⁷</p> |
| | <p>90% of tuna captured within 150 nm of tagging location.</p> | <p>Clear <i>et al.</i> (2005)¹⁰¹</p> |
| Albacore tuna | <p>Two individuals tagged and recaptured: one moved 302 nm, the other 1,727 nm.</p> | <p>I&I NSW (2009)⁹⁸</p> |
| | <p>Average distance travelled was 859.25</p> | <p>Cosgrove <i>et al.</i> (2010)¹⁰²</p> |

| Species | Info | Reference |
|----------------------------|--|--|
| | km. | |
| Dolphinfish | Move distances of up to 440 km. | Kingsford and Defries (1999) ¹⁰³ |
| | Average distance travelled for all individuals recaptured from 2006 to 2008 was 112.6 nm. | I&I NSW (2009) ⁹⁸ |
| Striped marlin | 90% of 360 tagged individuals were recaptured less than 1,000 km from the tagging location. | Jensen <i>et al.</i> (2010) ³² |
| | Mean straight-line distance per fish was 921 ± 264 km. | Holdsworth <i>et al.</i> (2009) ¹⁰⁴ |
| | Average distance of 280 nm | Squire Jr. (1974) ¹⁰⁵ |
| | The majority of striped marlin released off Australia have a mean displacement of less than 200 nm (after six to nine months). | Bromhead <i>et al.</i> (2004) ¹⁰⁶ |
| | Average distance travelled for all individuals recaptured from 2006 to 2008 was 214.2 nm. | I&I NSW (2009) ⁹⁸ |
| Shortfin mako shark | Move between ocean basins, enough to cause a lack of genetic differentiation. | Schrey and Heist (2003) ¹⁰⁷ |
| | Approximately 75% of the makos travelled less than 500 nm from their original tagging location with a mean distance of 398. | Kohler <i>et al.</i> (2002) ¹⁰⁸ |
| | Average distance for seven juveniles tracked between six and 45 hours was 55 km. | Sepulveda <i>et al.</i> (2004) ¹⁰⁹ |
| | Average distance travelled for all individuals recaptured from 2006 to 2008 was 571 nm. | I&I NSW (2009) ⁹⁸ |
| | Tagged off eastern Australia, stayed within the region. | Stevens <i>et al.</i> (2010) ¹¹⁰ |
| Blue shark | 82% of recaptured blue sharks travelled less than 1,000 km. | Queiroz <i>et al.</i> (2005) ¹¹¹ |
| | More than 75% of the blue sharks travelled less than 1,000 nm from their original tagging location with a | Kohler <i>et al.</i> (2002) ¹⁰⁸ |

| Species | Info | Reference |
|------------------------|--|---|
| | mean distance of 463 nm. | |
| | Average distance travelled for all individuals recaptured from 2006 to 2008 was 697 nm. | I&I NSW (2009) ⁹⁸ |
| | Tagged off eastern Australia, stayed within the region. | Stevens <i>et al.</i> (2010) ¹¹⁰ |
| Wahoo | Can move more than 1,000 km. | Theisen <i>et al.</i> (2008) ¹¹² |
| Skipjack tuna | Median lifetime displacement ranges from 420 to 470 nm. | Sibert and Hampton (2003) ⁹⁴ |
| Black marlin | Average short-term movement of five tagged marlin was 277.4 nm. | Gunn <i>et al.</i> (2003) ¹¹³ |
| | Average distance travelled for all individuals recaptured from 2006 to 2008 was 727.5 nm. | I&I NSW (2009) ⁹⁸ |
| Sailfish | Average distance travelled for all individuals recaptured from 2006 to 2008 was 38.6 nm. | I&I NSW (2009) ⁹⁸ |
| Porbeagle shark | More than 90% of tagged porbeagles travelled less than 500 nm from their original tagging location, with a mean distance of 234. | Kohler <i>et al.</i> (2002) ¹⁰⁸ |
| Thresher shark | Tagged off eastern Australia, stayed within the region. | Stevens <i>et al.</i> (2010) ¹¹⁰ |

Argument 2: Fisheries losses and overexploitation

Fishers are typically opposed to the establishment of marine reserves that overlap with their fishing grounds because of the perceived loss of revenue^{29,114}, even though it has been shown that fishing yield can be equivalent between marine reserves and traditional fisheries management¹¹⁵. Closure of large areas to fishing may lead to displacement of fishing effort that will cause overexploitation elsewhere^{3,31}. For instance, in a modelling study of the northwest Atlantic, the closure of an area with low to intermediate existing fishing effort simply displaced fishing effort elsewhere—to areas of higher species diversity—and increased the catch rate of most shark species³.

An economic impact assessment of fisheries closure in the high seas concluded that overall losses would be minimal in comparison with the ecosystem service and biodiversity benefits gained from such protection—it was calculated that the closure of 20 percent of the high seas may lead to the loss of only 1.8 percent of the current global reported marine fisheries catch¹¹⁶. It has now been established that marine reserves can export target species into adjacent fisheries, both through the emigration of adults and juveniles and the export of

propagules^{23,117, 118,119}. This “spillover” effect must be integrated into analyses of displaced fishing effort.

It is unclear whether overexploitation of pelagic species is occurring in the Coral Sea. Global assessments consider the region lightly fished⁶⁵, but estimates of catches on the northeast shelf, which include the Coral Sea, have documented large declines in catch rates (Figure 1). Furthermore, a number of species are considered at high risk from Eastern Tuna and Billfish Fishery (ETBF) longlining, most notably longfin mako sharks, crocodile sharks, pelagic thresher sharks, dusky sharks, ocean sunfish, short-finned pilot whales, false killer whales and leatherback turtles⁷⁶. Two commercial fisheries (the Coral Sea Fishery, or CSF, and the ETBF)ⁱ make up the bulk of the catch in the entire Coral Sea, and the charter fishing industry targets large predators in the Great Barrier Reef Marine Park and Coral Sea¹²⁰. The CSF is small (with 17 licensed operators, a mean gross value of production [GVP] of A\$866,000 between 2002 and 2006) and data-poor, making it difficult to ascertain trends and sustainability^{121,122}. The lack of good data will make it extremely difficult to calculate the impact of the proposed closure of the CSF¹²¹.

The ETBF also operates within the entire Coral Sea area but extends farther south, and less than one-third of its catch (between 15 and 31 percent in the years 1998 to 2006) comes from the Coral Sea itself (Table 3). The largest catch per unit effort for all species within the Coral Sea comes from its southern edge¹²³. Climate change is expected to drive many large pelagic species, including the 14 top species caught by the ETBF, farther south, suggesting that yields for this fishery in the Coral Sea are expected to decline⁹⁶.

The potential costs and displaced fishing effort of recreational and charter fishing in the Coral Sea is less well-understood. The sport fishery between Cairns and Lizard Island is highly lucrative, targeting a well-known black marlin spawning aggregation⁶⁰. Game fishing from charter vessels occurs around many reefs and seamounts of the Coral Sea, but data from the long-term Game Fish Tagging Program indicates that between 1989 and 2009, 99.2 percent of more than 18,000 tagged fish were caught in the Great Barrier Reef Marine Park and 0.8 percent in the Coral Sea¹²⁴. It is therefore possible that the risk of displacing fishing effort is small, and that populations of tuna, shark and billfish species that are heavily exploited outside the Coral Sea may benefit from marine reserve protection along this part of their range.

ⁱ Five fisheries are permitted to operate in the Coral Sea. Additional fisheries to the CSF and the ETBF are the Norfolk Island Fishery, the Southern Bluefin Tuna Fishery, the Eastern Skipjack Fishery and the Torres Strait Island Fishery. Catches from these fisheries are either almost entirely from outside the Coral Sea or the data are absorbed within the ETBF for reporting purposes.




Figure 1. Landings by functional groups on the northeast Australian shelf, which includes the Coral Sea, 1950-2006. From www.searoundus.org/lme/40/3.aspx.

Table 3. Catches (tonnes) of the most important ETBF species between 1998 and 2006. Data are split between catches from 25°S north (lying mostly in the Coral Sea), total catch from all areas, and the percent of the catch from above 25°S (%CS) from ¹²³.

| Species | Location | 98/99 | 99/00 | 00/01 | 01/02 | 02/03 | 03/04 | 04/05 | 05/06 |
|----------------------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|
| Yellowfin tuna | Coral Sea | 753 | 258 | 652 | 778 | 856 | 465 | 370 | 463 |
| | Total | 2,144 | 1,306 | 1,499 | 2,460 | 3,390 | 2,407 | 1,945 | 1523 |
| | % CS | 35.1 | 19.7 | 43.5 | 31.6 | 25.2 | 19.3 | 19 | 30.4 |
| Bigeye tuna | Coral Sea | 421 | 196 | 472 | 324 | 338 | 200 | 268 | 137 |
| | Total | 897 | 679 | 998 | 1,019 | 934 | 769 | 822 | 555 |
| | % CS | 46.9 | 28.9 | 47.3 | 31.8 | 36.2 | 26 | 32.6 | 24.7 |
| Broadbill swordfish | Coral Sea | 366 | 216 | 404 | 324 | 211 | 121 | 148 | 127 |
| | Total | 1,651 | 2,081 | 1,854 | 2,336 | 2,175 | 1,669 | 1,637 | 1,447 |
| | % CS | 22.2 | 10.4 | 21.8 | 13.9 | 9.7 | 7.2 | 9 | 8.8 |
| Striped marlin | Coral Sea | 50 | 36 | 138 | 150 | 90 | 48 | 23 | 80 |
| | Total | 492 | 514 | 717 | 768 | 631 | 574 | 408 | 505 |
| | % CS | 10.2 | 7 | 19.2 | 19.5 | 14.3 | 8.4 | 5.6 | 15.8 |
| Albacore | Coral Sea | 93 | 76 | 91 | 159 | 107 | 64 | 83 | 809 |

| Species | Location | 98/99 | 99/00 | 00/01 | 01/02 | 02/03 | 03/04 | 04/05 | 05/06 |
|---------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|
| | Total | 404 | 362 | 396 | 663 | 493 | 546 | 620 | 1584 |
| | % CS | 23 | 20.9 | 22.9 | 23.9 | 21.7 | 11.7 | 13.4 | 51.1 |
| Sharks | Coral Sea | 74 | 52 | 83 | 95 | 42 | 42 | 22 | 18 |
| | Total | 222 | 280 | 305 | 336 | 175 | 177 | 134 | 108 |
| | % CS | 33.3 | 18.6 | 27.2 | 28.3 | 24 | 23.7 | 16.4 | 16.7 |
| Other | Coral Sea | 126 | 45 | 209 | 219 | 129 | 128 | 100 | 143 |
| | Total | 334 | 267 | 862 | 756 | 644 | 657 | 546 | 602 |
| | % CS | 37.7 | 16.8 | 24.2 | 28.9 | 20 | 19.5 | 18.3 | 23.7 |
| TOTAL | Coral Sea | 1,884 | 880 | 2,050 | 2,051 | 1,773 | 1,068 | 1,014 | 1,777 |
| | Total | 6,143 | 5,488 | 6,631 | 8,338 | 8,442 | 6,798 | 6,111 | 6,324 |
| | % CS | 30.7 | 16 | 30.9 | 24.6 | 21 | 15.7 | 16.6 | 28.1 |

Argument 3: Governance and enforcement

The third key criticism of pelagic marine reserves centres on the costs and logistic constraints of governance, but the greatest concerns are directed toward marine reserves in international waters^{4,11,74}. Where a marine reserve is placed entirely within a nation's EEZ, concerns are focused less on governance issues and more on the logistics and costs of management¹¹. Large, multiple-use marine reserves such as the Great Barrier Reef Marine Park can offer both a blueprint for the type of surveillance and enforcement required, and a cautionary tale of the effects of insufficient enforcement activity^{10,125}.

There is still much to be learned about the enforcement of pelagic marine reserves¹²⁶. In remote oceanic areas, large no-take marine reserves are easier to police than multiuse parks or even marine reserve networks²², and a recent analysis found that larger reserves cost less per unit area than smaller ones¹²⁷. This same analysis examined three potential management scenarios for the Coral Sea (one large no-take reserve, a multiuse park with 30 percent of its area as no-take, and three smaller no-take areas making up 30 percent of the whole area together), and found that a single large no-take area would be the least costly to manage.

The proposed Coral Sea marine reserve would not extend beyond the boundaries of Australia's EEZ, precluding the need for complex international arrangements. Surveillance in this vast and remote area may be challenging, but there is increasing sophistication in vessel-monitoring systems (VMSs) and satellite technology¹¹. Additionally, there are existing arrangements and infrastructure that may be used for management activities and compliance monitoring¹²⁷ (see also Table 4). As a relatively wealthy nation with an extended EEZ, Australia is in an ideal position to take a leading role in the global protection of oceanic species and ecosystems.

Table 4. Summary table of challenges raised by critics of marine reserves for the protection of pelagic species, with proposed solutions and counterarguments, both general and directly relevant to the Coral Sea. Table reproduced and extended from Game *et al.* (2009) ¹¹.

| Issue | Challenges | Solutions and counter-arguments | Coral Sea example | |
|-------------------|--|---|--|--|
| Biological | Many pelagic species are highly mobile, often covering thousands of kilometres annually. | Spatial protection is either impossible across whole ranges or for all life-history stages, or the area required for conservation management would be unreasonably large. | Many threats to pelagic organisms are either site specific or cumulative, and can be reduced through spatial protection ³⁴ . In addition, many organisms either show site fidelity or have relatively small and defined areas of critical habitat within their range or life histories ^{38,58} . | The Coral Sea marine reserve would protect around 1 million km ² of ocean, including islands, reefs, seamounts, open ocean, abyssal plains, plateaux and canyons. Even highly mobile species have a high probability of spending at least 50% of their life cycle in the Coral Sea (Table 1). |
| | | Regulations or moratoria on gears or catch are more appropriate for limiting incidental capture of threatened pelagic fauna. | Although catch and gear regulations are an important component of pelagic conservation, they have so far proved inadequate in protecting many target and bycatch species ⁹⁵ . | The ETBF still records large quantities of bycatch, and stocks of its primary target species are known to be vulnerable ¹²⁷ . |
| Physical | The pelagic ocean is characterized by physical processes that are dynamic in space and time. | The environment is too dynamic to be represented in static reserves. Mobile reserves would be too difficult to enforce. | Many important pelagic features are either spatially or temporally predictable ⁸² , so static or dynamic marine reserves need to be designed accordingly ⁵⁵ . For features with less predictability, mobile fisheries closures have been effectively implemented off eastern Australia based on near real-time predictions | A Coral Sea marine reserve would encompass a number of permanent or predictable bathymetric and hydrographic features known to attract aggregations of pelagic and migratory species ⁶² . |

| | | | | |
|-------------------|---|--|--|--|
| Design | The pelagic ocean is generally data-poor compared with terrestrial or coastal systems. | Lack of data on the complexities of pelagic ecosystems limits the selection and design of marine reserves. | of pelagic habitat ⁵¹ . Governance issues are also addressed below. Widespread data sets, especially time-series data on remotely sensed physical and biological features (e.g., chlorophyll), are more abundant than commonly perceived and are useful for marine reserve selection. In contrast to fisheries catch limits, the selection of pelagic marine reserves does not have to rely on full understanding of ecosystem functions. | Remote sensing data for the Coral Sea is available through a number of sources. The general understanding of the Coral Sea's bathymetry (structure of the seafloor), oceanography and ecology is good ^{62,128} . |
| | There is a lack of well-established design principles to inform the selection of pelagic marine reserves. | Design principles for pelagic marine reserves will need to be developed de novo. | Some existing conservation planning tools and methods can be used in the pelagic ocean (e.g., Marxan), and good case studies are starting to appear. New challenges will lead to novel solutions with broad impact. | One of the primary requirements for the protection of pelagic and migratory species is that a marine reserve be large enough to encompass at least 50% of their home range ⁴⁴ . The Coral Sea marine reserve would protect around 1 million km ² . This area is highly likely to be adequate for the protection of most pelagic and migratory species that use its habitats. |
| Governance | Marine reserves might need to extend outside a country's EEZ. | Beyond national jurisdictions there is no legal basis for marine reserves. | Numerous existing international and regional agreements can be exercised to regulate marine reserves in the high seas ²⁷ . | The proposed Coral Sea marine reserve does not extend beyond Australia's EEZ ³⁰ . |

| | | | |
|--|---|--|--|
| Exploitation of the pelagic ocean is generally difficult and expensive to observe, and it is therefore challenging to enforce regulations. | Marine reserves will be more difficult and expensive to enforce, especially in developing nations, than traditional catch or gear restrictions. | Widespread adoption of satellite VMSs, and financial support for this in developing nations, will improve remote surveillance. | Australia is a wealthy nation where technology and surveillance operations are, to some extent, already in place ¹²⁹ . A precedent is in place for the use of existing operations for research and compliance monitoring ¹²⁷ . |
|--|---|--|--|

Parallels with Chagos/BIOT

The newest addition to a growing list of very large marine reserves is the Chagos Archipelago or British Indian Ocean Territory, referred to by Koldewey *et al.* (2010)²⁷ as Chagos/BIOT. The ecological rationale behind the establishment of this marine reserve could easily be applied to the proposed Coral Sea marine reserve. Scientific research suggests that with 544,000 km² of ocean, reefs and seamounts, the Chagos/BIOT Marine Reserve potentially provides an excellent area for the recovery of shark, tuna and other large predators²⁷. Similarly, the Coral Sea is highly likely to be large enough (approximately 1 million km²) to adequately protect large pelagic predators that are subject to overexploitation elsewhere along their range¹³⁰. Other similarities between the two regions are the relatively intact nature of the entire system, the remoteness of the coral reefs from human impacts, their status as stepping-stones to genetic connectivity between broader regions, their regional importance for threatened and endangered species and the diversity of undersea geological features²⁷. These criteria were crucial in driving the protection of Chagos/BIOT⁴⁸.

Chagos/BIOT has met with opposition, especially from the tuna fishing industry²². Unfortunately, the lack of data about tuna catches within the reserve made it difficult to estimate the potential loss to the industry or the displacement likely to occur into other areas^{22,48}. Estimates of the proportion of the Indian Ocean catch coming from the Chagos/BIOT area range from 2 to 6 percent²². The proportion of the ETBF catch in the Coral Sea is higher, ranging from 15 to 31 percent between 1998 and 2006. The Coral Sea Fishery operates entirely within the proposed reserve area, but its commercial value is relatively low. Recreational fishers and representatives of the game-fishing industry, rather than the commercial fishing industry, have expressed opposition to the protection of the Coral Sea (I. Zethoven, pers. comm.).

Chagos/BIOT was declared in the first half of 2010 despite a lack of existing information, apart from some fisheries data²⁷. In comparison, despite knowledge gaps, the Coral Sea is reasonably well-understood; knowledge of underlying bathymetric and oceanographic features and biological communities has allowed broad ecological patterns and processes to be described or inferred⁶². There is increasing global demand to heed the precautionary principle, especially in marine ecosystems where data are scarce and fisheries decline or collapse has been well-documented^{13,43,44,82}.

Conclusions

This review has detected a trend in the scientific literature toward greater support for the use of marine reserves to protect migratory and highly mobile pelagic species. Recent research has demonstrated that large pelagic species targeted by fisheries benefit from marine reserves. This is most likely because:

- Even protecting a part of species' ranges or life cycles, especially critical habitat areas which function as important feeding or breeding grounds, can cause a decrease in overall population mortality;
- Pelagic species are not uniformly distributed, but tend to aggregate around bathymetric and hydrographic features that are predictable in space and time, making the most beneficial design of pelagic reserves possible; and

- Even species identified as “highly migratory” display movement patterns where the majority (70 to 90 percent) of the population moves no farther than 600 km.

Critics of marine reserves for the conservation of pelagic species have focused on three major issues. Firstly, there has been a concern that reserves have to be very large to encompass an adequate portion of pelagic species’ home ranges. Secondly, there is some scepticism about the use of marine reserves as a fisheries management tool, with concerns about displaced fishing effort causing overexploitation in adjacent areas. Thirdly, there is a great degree of difficulty associated with arranging governance and enforcement beyond one country’s EEZ .

The proposed Coral Sea marine reserve is ideally placed to surmount the challenges raised. The proposed reserve encompasses almost 1 million km², making it large enough to protect a significant portion (if not all) of the home ranges and life cycles of most pelagic species that reside within it. It is large enough to also encompass a large variety of bathymetric and hydrographic features that provide key habitat for pelagic species at vulnerable times (feeding and breeding). It lies entirely within Australia’s EEZ, making difficult negotiations with neighbouring countries unnecessary. Existing marine operations and technology may be used for surveillance and enforcement. The Coral Sea marine reserve is being proposed as a conservation tool rather than a fisheries management strategy. Its history of relatively low exploitation, coupled with its relatively high remaining density and diversity of large pelagic predators, is considered ideal for conservation, as its ecosystems remain relatively undisturbed. Finally, a recent precedent exists: The ecological values leading to the declaration of the Chagos/BIOT marine reserve are also found in the Coral Sea.

Acknowledgments

This paper has benefited greatly from peer review by Dr. Maria Beger and Dr. Natalie Ban of the Australian Research Council’s Centre of Excellence for Coral Reef Studies and Dr. Edward Game of The Nature Conservancy.

References

1. Verity, P.G., *et al.* (2002) Status, trends and the future of the marine pelagic ecosystem. *Environmental Conservation* 29, 207-237
2. Mills, C.E., and Carlton, J.T. (1998) Rationale for a system of international reserves for the open ocean. *Conservation Biology* 12, 244-247
3. Baum, J.K., *et al.* (2003) Collapse and conservation of shark populations in the Northwest Atlantic. *Science* 299, 389-392
4. de Juan, S., and Leonart, J. (2010) A conceptual framework for the protection of vulnerable habitats impacted by fishing activities in the Mediterranean high seas. *Ocean and Coastal Management* 53, 717-723
5. Boersma, P.D., and Parrish, J.K. (1999) Limiting abuse: marine protected areas, a limited solution. *Ecological Economics* 31, 287-304
6. Myers, R.A., *et al.* (2007) Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science* 315, 1846-1850
7. Graham, N.A.J., *et al.* (2010) Reef shark declines in remote atolls highlight the need for multi-faceted conservation action. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20, 543-548
8. Ward-Paige, C.A., *et al.* (2010) Large-scale absence of sharks on reefs in the greater-Caribbean: A footprint of human pressures. *PLoS ONE* 5(8): e11968. doi:10.1371/journal.pone.0011968
9. Wood, L.J., *et al.* (2008) Assessing progress towards global marine protection targets: shortfalls in information and action. *Oryx* 42, 340-351
10. McCook, L.J., *et al.* (2010) Adaptive management of the Great Barrier Reef: A globally significant demonstration of the benefits of networks of marine reserves. *PNAS*
www.pnas.org/cgi/doi/10.1073/pnas.0909335107
11. Game, E.T., *et al.* (2009) Pelagic protected areas: the missing dimension in ocean conservation. *Trends in Ecology and Evolution* 24, 360-369
12. UNEP (2010) COP 10 Decision X/2 - Strategic Plan for Biodiversity 2011-2020.
<http://www.cbd.int/decision/cop/?id=12268>

13. Foley, M.M., *et al.* (2010) Guiding ecological principles for marine spatial planning. *Marine Policy* 34, 955-966
14. Caley, M.J., and St John, J. (1996) Refuge availability structures assemblages of tropical reef fishes. *Journal of Animal Ecology* 65, 414-428
15. Graham, N.A.J., *et al.* (2009) Coral mortality versus structural collapse as drivers of corallivorous butterflyfish decline. *Biodiversity and Conservation* 18, 3325-3336
16. Jones, G.P., *et al.* (2007) Population connectivity and conservation of marine biodiversity. *Oceanography* 20, 100-111
17. Worm, B., *et al.* (2006) Impacts of biodiversity loss on ocean ecosystem services. *Science* 314, 787-790
18. Palumbi, S.R., *et al.* (2008) Ecosystems in action: lessons from marine ecology about recovery, resistance, and reversibility. *BioScience* 58, 33-42
19. Dulvy, N.K., *et al.* (2004) Coral reef cascades and indirect effects of predator removal by exploitation. *Ecology Letters* 7, 410-416
20. Bellwood, D.R., *et al.* (2004) Confronting the coral reef crisis. *Nature* 429, 827-833
21. Berke, S.K. (2010) Functional groups of ecosystem engineers: A proposed classification with comments on current issues. *Integrative and Comparative Biology* 50, 147-157
22. Sheppard, C. (2010) Marine protected areas and pelagic fishing: The case of the Chagos Archipelago. *Marine Pollution Bulletin* 60, 1899-1901
23. Gell, F.R., and Roberts, C.M. (2003) Benefits beyond boundaries: the fishery effects of marine reserves. *Trends in Ecology and Evolution* 18, 448-455
24. Halpern, B.S., and Warner, R.R. (2002) Marine reserves have rapid and lasting effects. *Ecology Letters* 5, 361-366
25. Claudet, J., *et al.* (2008) Marine reserves: size and age do matter. *Ecology Letters* 11, 481-489
26. Willis, T.J., *et al.* (2003) Protection of exploited fish in temperate regions: high density and biomass of snapper *Pagrus auratus* (Sparidae) in northern New Zealand marine reserves. *Journal of Applied Ecology* 40, 214-227

27. Koldewey, H.J., *et al.* (2010) Potential benefits to fisheries and biodiversity of the Chagos Archipelago/British Indian Ocean Territory as a no-take marine reserve. *Marine Pollution Bulletin* 60, 1906-1915
28. Micheli, F., *et al.* (2004) Trajectories and correlates of community change in no-take marine reserves. *Ecological Applications* 14, 1709-1723
29. Nelson, J., and Bradner, H. (2010) The case for establishing ecosystem-scale marine reserves. *Marine Pollution Bulletin* 60, 635-637
30. Zethoven, I.e. (2008) An Australian Coral Sea heritage park., The Pew Environment Group
31. Worm, B., *et al.* (2003) Predator diversity hotspots in the blue ocean. *PNAS* 100, 9884–9888
32. Jensen, O.P., *et al.* (2010) Local management of a “highly migratory species”: The effects of long-line closures and recreational catch-and-release for Baja California striped marlin fisheries. *Progress in Oceanography* 86, 176-186
33. Claudet, J., *et al.* (2010) Marine reserves: fish life history and ecological traits matter. *Ecological Applications* 20, 830-839
34. Hooker, S.K., and Gerber, L.R. (2004) Marine reserves as a tool for ecosystem-based management: The potential importance of megafauna. *BioScience* 54, 27-39
35. Beare, D., *et al.* (2010) An unintended experiment in fisheries science: a marine area protected by war results in Mexican waves in fish numbers-at-age. *Naturwissenschaften* 97, 797-808
36. Roberts, R.C., and Sargant, H. (2002) Fishery benefits of fully protected marine reserves: why habitat and behaviour are important. *Natural Resource Modeling* 15, 487-507
37. Palumbi, S.R. (2004) Marine reserves and ocean neighborhoods: The spatial scale of marine populations and their management. *Annual Review of Environment and Resources* 29, 31-68
38. Louzao, M., *et al.* (2006) Oceanographic habitat of an endangered Mediterranean porcellariiform: Implications for marine protected areas. *Ecological Applications* 16, 1683-1695
39. Pichegru, L., *et al.* (2009) Overlap between vulnerable top predators and fisheries in the Benguela upwelling system: implications for marine protected areas. *Marine Ecology Progress Series* 391, 199-208

40. Gibson, L.E., and Wellbelove, A.P. (2010) Protecting Critical Marine Habitats: The key to conserving our threatened marine species., A Humane Society International and WWF-Australia Report
41. Claydon, J. (2004) Spawning aggregations of coral reef fishes: Characteristics, hypotheses, threats and management. *Oceanography and Marine Biology: an Annual Review* 42, 265-302
42. Pears, R.J., *et al.* (2006) Demography of a large grouper, *Epinephelus fuscoguttatus*, from Australia's Great Barrier Reef: implications for fishery management. *Marine Ecology Progress Series* 307, 259-272
43. Lauck, T., *et al.* (1998) Implementing the precautionary principle in fisheries management through marine reserves. *Ecological Applications* 8, S72-S78
44. Clark, C.W. (1996) Marine reserves and the precautionary management of fisheries. *Ecological Applications* 6, 369-370
45. Gerber, L.R., *et al.* (2005) Do the largest protected areas conserve whales or whalers? *Science* 307, 525-526
46. Walters, C. (2000) Impacts of dispersal, ecological interactions, and fishing effort dynamics on efficacy of marine protected areas: How large should protected areas be? *Bulletin of Marine Science* 66, 645-757
47. Griffiths, M.H., and Wilke, C.G. (2002) Long-term movement patterns of five temperate-reef fishes (Pisces:Sparidae): implications for marine reserves. *Marine and Freshwater Research* 53, 233-244
48. Williamson, P. (2010) Marine conservation in the British Indian Ocean Territory: science issues and opportunities. Workshop held 5-6 August 2009 at the National Oceanography Centre, Southampton supported by the NERC Strategic Ocean Funding Initiative and the Oceans 2025 programme., University of East Anglia Print Services
49. Hilborn, R., *et al.* (2004) When can marine reserves improve fisheries management? *Ocean and Coastal Management* 47, 197-205
50. Corrigan, C., and Kershaw, F. (2008) Working toward high seas marine protected areas: An assessment of progress made and recommendations for collaboration., UNEP-WCMC
51. Hobday, A.J., and Hartmann, K. (2006) Near real-time spatial management based on habitat predictions for a longline bycatch species. *Fisheries Management and Ecology* 13, 365-380

52. Notarbatolo di Sciara, G., *et al.* (2008) The Pelagos Sanctuary for Mediterranean marine mammals. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18, 367-391
53. Burnell, S.R., and Bryden, M.M. (1997) Coastal residence periods and reproductive timing in southern right whales, *Eubalaena australis*. *Journal of Zoology, London* 241, 613-621
54. Pala, C. (2009) Protecting the last great tuna stocks. *Science* 324, 1133
55. Alpine, J.E., and Hobday, A.J. (2007) Area requirements and pelagic protected areas: is size an impediment to implementation? *Marine and Freshwater Research* 58, 558-569
56. Richards, R. (2009) Past and present distributions of southern right whales (*Eubalaena australis*). *New Zealand Journal of Zoology* 36, 447-459
57. Clapham, P.J., *et al.* (1999) Baleen whales: conservation issues and the status of the most endangered populations. *Mammal Review* 29, 35-60
58. Hyrenbach, K.D., *et al.* (2006) Use of marine sanctuaries by far-ranging predators: commuting flights to the California Current System by breeding Hawaiian albatrosses. *Fisheries Oceanography* 15, 95-103
59. Young, J., *et al.* (2003) Reproductive dynamics of broadbill swordfish, *Xiphias gladius*, in the domestic longline fishery off eastern Australia. *Marine and Freshwater Research* 54, 315-332
60. Speare, P. (2003) Age and growth of black marlin, *Makaira indica*, in east coast Australian waters. *Marine and Freshwater Research* 54, 307-314
61. Farley, J.H., *et al.* (2004) Age and growth of bigeye tuna (*Thunnus obesus*) in the eastern and western AFZ. *IOTC-2004-WPTT-INF04*
62. Brewer, D.T., *et al.* (2007) Ecosystems of the East Marine Planning Region., Report to Department of Environment and Water Resources. CSIRO
63. Moritz, C., *et al.* (2002) Population genetics of Southeast Asian and Western Pacific green turtles, *Chelonia mydas*., Reports to UNEP / CMS
64. Congdon, B.C., *et al.* (2007) Vulnerability of seabirds on the Great Barrier Reef to climate change. In *Climate Change and the Great Barrier Reef: A Vulnerability Assessment*. (Johnson, J.E., and Marshall, P.A., eds), Great Barrier Reef Marine Park Authority

65. Mora, C., *et al.* (2009) Management effectiveness of the world's marine fisheries. *PLoS Biology* 7: e1000131. doi:10.1371/journal.pbio.1000131
66. Baker, B., *et al.* (2008) Seabird monitoring study at Coringa Herald National Nature Reserve., Report to the Department of the Environment, Water, Heritage and the Arts by Latitude 42 Environmental Consultants Pty Ltd
67. Birtles, A., *et al.* (2008) Marine and Tropical Sciences Research Facility Annual Report, June 2008. Program 8: Sustainable Use and Management of Marine Resources of the Great Barrier Reef., Marine and Tropical Sciences Research Facility
68. Kaiser, M.J. (2005) Are marine protected areas a red herring or fisheries panacea? *Canadian Journal of Fisheries and Aquatic Sciences* 62, 1194-1199
69. Lubchenco, J., *et al.* (2003) Plugging a hole in the ocean: The emerging science of marine reserves. *Ecological Applications* 13, S3-S7
70. Hughes, T.P., *et al.* (2010) Rising to the challenge of sustaining coral reef resilience. *Trends in Ecology and Evolution* 25, 633-642
71. Williams, A., *et al.* (2010) Seamount megabenthic assemblages fail to recover from trawling impacts. *Marine Ecology* 31, 183-199
72. Hughes, T.P., *et al.* (2003) Climate change, human impacts, and the resilience of coral reefs. *Science* 301, 929-933
73. Norse, E.A. (2005) Pelagic protected areas: the greatest parks challenge of the 21st century. *Parks* 15:32-39. *Parks* 15, 32-39
74. Kaplan, D.M., *et al.* (2010) Pelagic MPAs: The devil is in the details. *Trends in Ecology and Evolution* 25, 62-63
75. Davies, A.J., *et al.* (2007) Preserving deep-sea natural heritage: Emerging issues in offshore conservation and management. *Biological Conservation* 138, 299-312
76. Wilson, D.T., *et al.*, eds (2010) *Fishery status reports 2009: status of fish stocks and fisheries managed by the Australian Government*. Australian Bureau of Resource Economics - Bureau of Rural Sciences
77. Ceccarelli, D., *et al.* (2009) Lihou Reef National Nature Reserve Marine Survey – October 2008., Report to the Department of the Environment, Water, Heritage and the Arts by C&R Consulting Pty Ltd.

78. Planes, S., *et al.* (2001) Strong genetic divergence among populations of a marine fish with limited dispersal, *Acanthochromis polycanthus*, within the Great Barrier Reef and the Coral Sea. *Evolution* 55, 2263-2273
79. Benzie, J.A.H., and Williams, S.T. (1992) Genetic structure of giant clam (*Tridacna maxima*) populations from reefs in the Western Coral Sea. *Coral Reefs*, 11(3): 135-141. *Coral Reefs* 11, 135-141
80. van Herwerden, L., *et al.* (2009) Complex patterns of population structure and recruitment of *Plectropomus leopardus* (Pisces: Epinephelidae) in the Indo-West Pacific: implications for fisheries management. *Marine Biology* 156, 1595-1607
81. Sarano, F., and Pichon, M. (1988) Morphology and ecology of the deep fore reef slope at Osprey Reef (Coral Sea). In *Proceedings of the 6th International Coral Reef Symposium*, 607-611
82. Hyrenbach, K.D., *et al.* (2000) Marine protected areas and ocean basin management. *Aquatic Conservation: Marine and Freshwater Ecosystems* 10, 437-458
83. Jorgensen, S.J., *et al.* (2009) Philopatry and migration of Pacific white sharks. *Proceedings of the Royal Society B: Biological Sciences* doi: 10.1098/rspb.2009.1155
84. Rissik, D., and Suthers, I.M. (2000) Enhanced feeding by pelagic juvenile myctophid fishes within a region of island-induced flow disturbance in the Coral Sea. *Marine Ecology Progress Series* 203, 263-273
85. Morato, T., *et al.* (2008) Evidence of a seamount effect on aggregating visitors. *Marine Ecology Progress Series* 357, 23-32
86. Hyrenbach, K., *et al.* (2002) Oceanographic habitats of two sympatric North Pacific albatrosses during breeding season. *Marine Ecology Progress Series* 233, 283-301
87. Hearn, A., *et al.* (2010) Hotspots within hotspots? Hammerhead shark movements around Wolf Island, Galapagos Marine Reserve. *Marine Biology* 157, 1899-1915
88. Etnoyer, P., *et al.* (2006) Sea-surface temperature gradients across blue whale and sea turtle foraging trajectories off the Baja California Peninsula, Mexico. *Deep-Sea Research II* 53, 340-358
89. Ferraroli, S., *et al.* (2004) Where leatherback turtles meet fisheries. *Nature* 429, 521
90. Keene, J., *et al.* (2008) Sedimentology and geomorphology of the East Marine region of Australia., Report to the Department of the Environment, Water, Heritage and the Arts by Geoscience Australia

91. Choukroun, S., *et al.* (2010) On the surface circulation in the western Coral Sea and residence times in the Great Barrier Reef. *Journal of Geophysical Research* 115, C06013, doi:06010.01029/02009JC005761
92. Fowler, A.M., *et al.* (2008) Onshore-offshore distribution and abundance of tuna larvae (Pisces: Scombridae: Thunnini) in near-reef waters of the Coral Sea. *Fisheries Bulletin* 106, 405-416
93. Bradshaw, C.J.A., *et al.* (2008) Decline in whale shark size and abundance at Ningaloo Reef over the past decade: The world's largest fish is getting smaller. *Biological Conservation* 141, 1894-1905
94. Sibert, J., and Hampton, J. (2003) Mobility of tropical tunas and the implications for fisheries management. *Marine Policy* 27, 87-95
95. Pauly, D., *et al.* (2002) Towards sustainability in world fisheries. *Nature* 418, 689-695
96. Hobday, A.J. (2010) Ensemble analysis of the future distribution of large pelagic fishes off Australia. *Progress in Oceanography* 86, 291-301
97. Hampton, J., and Gunn, J. (1998) Exploitation and movements of yellowfin tuna (*Thunnus albacares*) and bigeye tuna (*T. obesus*) tagged in the north-western Coral Sea. *Marine and Freshwater Research* 49, 475-489
98. I&I NSW (2009) I&I NSW Game Fish Tagging Program, Summary 2006/2007 & 2007/2008., I&I New South Wales
99. Sedberry, G.R., and Loefer, J.K. (2001) Satellite telemetry tracking of swordfish, *Xiphias gladius*, off the eastern United States. *Marine Biology* 139, 355-360
100. Sepulveda, C.A., *et al.* (2010) Fine-scale movements of the swordfish *Xiphias gladius* in the Southern California Bight. *Fisheries Oceanography* 19, 279-289
101. Clear, N.P., *et al.* (2005) Movement of bigeye tuna (*Thunnus obesus*) determined from archival tag light-levels and sea surface temperatures., CSIRO, Hobart, and SPC
102. Cosgrove, R., *et al.* (2010) A simulated archival tagging programme for albacore (*Thunnus alalunga*) in the Northeast Atlantic, including an analysis of factors affecting tag recovery. *ICES Journal of Marine Science* 67, 1216-1221
103. Kingsford, M.J., and Defries, A. (1999) The ecology of and fishery for *Coryphaena* spp. in the waters around Australia and New Zealand. *Scientia Marina* 63, 267-275

104. Holdsworth, J.C., *et al.* (2009) Near real time satellite tracking of striped marlin (*Kajikia audax*) movements in the Pacific Ocean. *Marine Biology* 156, 505-514
105. Squire Jr., J.L. (1974) Migration patterns of Istiophoridae in the Pacific Ocean as determined by cooperative tagging programs., NOAA
106. Bromhead, D., *et al.* (2004) Striped marlin: biology and fisheries., Bureau of Rural Sciences
107. Schrey, A.W., and Heist, E.J. (2003) Microsatellite analysis of population structure in the shortfin mako (*Isurus oxyrinchus*). *Canadian Journal of Fisheries and Aquatic Sciences* 60, 670-675
108. Kohler, N.E., *et al.* (2002) Tag and recapture data for three pelagic shark species: Blue shark (*Prionace glauca*), shortfin mako (*Isurus oxyrinchus*), and porbeagle (*Lamna nasus*) in the north Atlantic Ocean. *Col. Vol. Sci. Pap. ICCAT* 54, 1231-1260
109. Sepulveda, C.A., *et al.* (2004) Movement patterns, depth preferences, and stomach temperatures of free-swimming juvenile mako sharks, *Isurus oxyrinchus*, in the Southern California Bight. *Marine Biology* 145, 191-199
110. Stevens, J.D., *et al.* (2010) Satellite tagging of blue sharks (*Prionace glauca*) and other pelagic sharks oV eastern Australia: depth behaviour, temperature experience and movements. *Marine Biology* 157, 575-591
111. Queiroz, N., *et al.* (2005) Movement of blue shark, *Prionace glauca*, in the north-east Atlantic based on mark-recapture data. *Journal of the Marine Biological Associations of the United Kingdom* 85, 1107-1112
112. Theisen, T.C., *et al.* (2008) High connectivity on a global scale in the pelagic wahoo, *Acanthocybium solandri* (tuna family Scombridae). *Molecular Ecology* 17, 4233-4247
113. Gunn, J.S., *et al.* (2003) Short-term movement and behaviour of black marlin *Makaira indica* in the Coral Sea as determined through a pop-up satellite archival tagging experiment. *Marine and Freshwater Research* 54, 515-525
114. Blyth-Skyrme, R.E., *et al.* (2006) Conservation benefits of temperate marine protected areas: variation among fish species. *Conservation Biology* 20, 811-820
115. Hastings, A., and Botsford, L.W. (1999) Equivalence in yield from marine reserves and traditional fisheries management. *Science* 284, 1537-1538
116. Sumaila, U.R., *et al.* (2007) Potential costs and benefits of marine reserves in the high seas. *Marine Ecology Progress Series* 345, 305-310

117. Goñi, R., *et al.* (2010) Net contribution of spillover from a marine reserve to fishery catches. *Marine Ecology Progress Series* 400, 233-243
118. Roberts, C.M., *et al.* (2001) Effects of marine reserves on adjacent fisheries. *Science* 294, 1920-1923
119. Halpern, B.S., *et al.* (2009) Spillover from marine reserves and the replenishment of fished stocks. *Environmental Conservation* 36, 268-276
120. Wilson, D., *et al.*, eds (2009) *Fishery status reports 2008: status of fish stocks and fisheries managed by the Australian Government*. Bureau of Rural Sciences & Australian Bureau of Agricultural and Resource Economics
121. Dowling, N.A., *et al.* (2008) Developing harvest strategies for low-value and data-poor fisheries: Case studies from three Australian fisheries. *Fisheries Research* 94, 380-390
122. Furlani, D., *et al.* (2007) Ecological risk assessment for the effects of fishing: Report for the demersal trawl sub-fishery of the Coral Sea Fishery., Report for the Australian Fisheries Management Authority
123. Evans, S.K. (2007) Eastern Tuna and Billfish Fishery Data Summary 2005-2006., Data Group, Australian Fisheries Management Authority
124. NSW DPI (2009) Game Fish Tagging Program.
125. Ayling, A.M., and Choat, J.H. (2008) Abundance patterns of reef sharks and predatory fishes on differently zones reefs in the offshore Townsville region., Report by Sea Research for the Great Barrier Reef Marine Park Authority
126. Game, E.T., *et al.* (2010) Pelagic MPAs: The devil you know. *Trends in Ecology and Evolution* 25, 63-64
127. Ban, N.C., *et al.* (2009) Marine protected area management costs: an analysis of options for the Coral Sea., Report prepared for the Protect Our Coral Sea campaign by the Australian Research Council Centre of Excellence for Coral Reef Studies
128. Beaman, R.J. (2010) 3DGBR: a high-resolution depth model for the Great Barrier Reef and Coral Sea. Marine and Tropical Sciences Research Facility Final Report Project 2.5i.1a Milestone 10 June 2010., Marine and Tropical Sciences Research Facility, Reef and Rainforest Research Centre
129. Brooke, S.D., *et al.* (2010) Surveillance and enforcement of remote maritime areas. Paper 1: surveillance technical options., Marine Conservation Biology Institute

130. Worm, B., *et al.* (2005) Global patterns of predator diversity in the open oceans. *Science* 309, 1365-1369

Appendix 1—Migratory and pelagic species of the Coral Sea

This table lists those migratory and pelagic species affected by fishing in the Coral Sea, either as target species or bycatch⁷⁶, and may therefore benefit from protection.

| Species | Common name | Pressure |
|---|------------------------|--------------------|
| <i>Thunnus albacares</i> | Yellowfin tuna | Commercial fishing |
| <i>Thunnus obesus</i> | Bigeye tuna | Commercial fishing |
| <i>Xiphias gladius</i> | Broadbill swordfish | Commercial fishing |
| <i>Thunnus alalunga</i> | Albacore tuna | Commercial fishing |
| <i>Coryphaena hippurus</i> | Dolphinfish | Commercial fishing |
| <i>Centrolophus niger</i> | Rudderfish | Commercial fishing |
| <i>Tetrapturus audax</i> | Striped marlin | Commercial fishing |
| <i>Isurus oxyrinchus</i> | Shortfin mako shark | Commercial fishing |
| <i>Prionace glauca</i> | Blue shark | Commercial fishing |
| <i>Acanthocybium solandri</i> | Wahoo | Commercial fishing |
| <i>Makaira indica</i> | Black marlin | Commercial fishing |
| <i>Brama brama</i> | Ray's bream | Commercial fishing |
| <i>Ruvettus pretiosus</i> | Black oilfish | Commercial fishing |
| <i>Katsuwonus pelamis</i> | Skipjack tuna | Commercial fishing |
| <i>Thunnus thynnus</i> | Northern bluefin tuna | Commercial fishing |
| <i>Tetrapturus angustirostris</i> | Shortbilled spearfish | Commercial fishing |
| <i>Istiophorus platypterus</i> | Indo-Pacific sailfish | Commercial fishing |
| <i>Lampris guttatus, L. immaculatus</i> | Moonfish | Commercial fishing |
| <i>Thunnus maccoyii</i> | Southern bluefin tuna | Commercial fishing |
| <i>Carcharhinus brachyurus</i> | Bronze whaler | Commercial fishing |
| <i>Carcharhinus longimanus</i> | Oceanic whitetip shark | Commercial fishing |

| | | |
|-----------------------------------|--------------------------------|--------------------|
| <i>Sphyrna spp.</i> | Hammerhead shark | Commercial fishing |
| <i>Galeocerdo cuvier</i> | Tiger shark | Commercial fishing |
| <i>Carcharhinus falciformis</i> | Silky shark | Commercial fishing |
| <i>Sarda australis</i> | Australian bonito | Commercial fishing |
| <i>Carcharhinus limbatus</i> | Dusky shark | Commercial fishing |
| <i>Lamna nasus</i> | Porbeagle | Commercial fishing |
| <i>Alopias vulpinus</i> | Thresher shark | Commercial fishing |
| <i>Scomberomorus commerson</i> | Narrow barred Spanish mackerel | Commercial fishing |
| <i>Rachycentron canadum</i> | Black kingfish | Commercial fishing |
| <i>Isurus paucus</i> | Longfin mako | Commercial fishing |
| <i>Carcharodon carcharias</i> | Great white shark | Bycatch |
| <i>Elegatis bipinnulata</i> | Rainbow runner | Bycatch |
| <i>Pseudocarcharias kamoharai</i> | Crocodile shark | Bycatch |
| <i>Caranx ignobilis</i> | Giant trevally | Bycatch |
| <i>Alepisaurus ferox</i> | Lancetfish | Bycatch |
| <i>Manta birostris</i> | Manta ray | Bycatch |
| Marine turtles | Various | Bycatch |
| Seabirds | Various | Bycatch |
